

Corrections to *ghm22@cam.ac.uk*. Star (★) indicates a harder question.

- 1 Use the Cauchy–Schwarz inequality and the properties of the inner product to prove the triangle inequality

$$|x + y| \leq |x| + |y|$$

for a complex vector space, where  $|x|$  is the norm of the vector  $x$ . Under what conditions does equality hold?

- 2 Given a set of vectors  $u_1, u_2, \dots, u_m$  ( $m \geq n$ ) that span an  $n$ -dimensional vector space, show that an orthogonal basis may be constructed by the Gram-Schmidt procedure:

$$e_1 = u_1$$

$$e_r = u_r - \sum_{s=1}^{r-1} \frac{e_s \cdot u_r}{e_s \cdot e_s} e_s \quad \text{for } r > 1.$$

What is the interpretation if any of the vectors  $e_r$  vanishes?

Find an orthonormal basis for the subspace of a four-dimensional Euclidean space spanned by the three vectors with components  $(1, 1, 0, 0)$ ,  $(0, 1, 2, 0)$  and  $(0, 0, 3, 4)$ .

- 3 An  $n \times n$  complex matrix  $A$  is such that each row and each column has exactly one non-zero element. The Hermitian conjugate of  $A$  is  $A^\dagger = (A^T)^*$  (where  $A^T$  is the transpose of  $A$ , and  $A^*$  is its complex conjugate). Show that  $A^\dagger A$  is a real diagonal matrix.
- 4★ What does it mean to say that the vectors  $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$  are linearly independent?

Let  $A$  be a linear operator on an  $n$ -dimensional vector space, having  $n$  distinct eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_n$  and corresponding eigenvectors  $e_1, e_2, \dots, e_n$ . Consider the action of the operator  $A - \lambda_i I$  (where  $I$  is the identity operator) on the vector  $e_j$  in the cases  $i = j$  and  $i \neq j$ .

Describe the action of this operator on  $\sum_j \alpha_j e_j$ . Hence or otherwise show that the vectors  $\{\mathbf{e}_i\}$  are linearly independent.

- 5 An  $n \times n$  Hermitian matrix  $A$  is said to be positive definite if  $\mathbf{x}^\dagger A \mathbf{x} > 0$  for all non-zero vectors  $\mathbf{x} \in \mathbb{C}^n$ .

Show that if  $A$  is positive definite, then any principal submatrix (a square matrix formed by deleting the same set of rows and columns from  $A$ ) is also positive definite. Deduce that all diagonal elements  $A_{ii}$  must be strictly positive real numbers.

For a  $2 \times 2$  positive definite matrix  $A$ , prove that  $\det A \leq A_{11}A_{22}$ . Under what condition does equality hold?

- 6 An Hermitian matrix  $A$  is one for which  $A^\dagger = A$ . Suppose that  $A$  and  $B$  are both Hermitian matrices. Show that  $AB + BA$  is Hermitian. Also show that  $AB$  is Hermitian if and only if  $A$  and  $B$  commute.

- 7 Suppose  $A$  is an  $n \times n$  matrix such that  $A^2 = A$ . By considering the eigenvalues of  $A$  show that either:

- $\det A = 1$  and  $\text{Tr } A = n$ ; or,
- $\det A = 0$  and  $\text{Tr } A = m$  with  $m < n$ .

Show that the matrix  $B = I - A$  also satisfies  $B^2 = B$ . Hence, show that any vector  $\mathbf{x}$  can be uniquely decomposed as  $\mathbf{x} = \mathbf{y} + \mathbf{z}$ , where  $A\mathbf{y} = \mathbf{y}$  and  $A\mathbf{z} = \mathbf{0}$ . What are the dimensions of the subspaces containing  $\mathbf{y}$  and  $\mathbf{z}$  in case (b)? Provide a geometric interpretation.

8 Show that two diagonalizable matrices  $A$  and  $B$  share a common basis of eigenvectors if and only if they commute. Use this result to deduce that the matrices  $(I + A)$  and  $(I - A)^{-1}$  commute.

9 An  $n \times n$  complex matrix  $N$  is said to be nilpotent if  $N^k = 0$  for some positive integer  $k$ .

By considering the eigenvalue equation  $N\mathbf{x} = \lambda\mathbf{x}$  for a non-zero eigenvector  $\mathbf{x}$ , show that the only eigenvalue of a nilpotent matrix is  $\lambda = 0$ . Hence, determine  $\det N$  and  $\text{Tr } N$ .

Show that the matrix  $(I - N)$  is always invertible, and find an explicit expression for  $(I - N)^{-1}$  as a finite sum involving powers of  $N$ .

10 In quantum mechanics, the statistical state of a system is often represented by a *density matrix*  $\rho$ . A valid density matrix must be Hermitian, positive semi-definite ( $\mathbf{x}^\dagger \rho \mathbf{x} \geq 0$  for all  $\mathbf{x}$ ), and satisfy  $\text{Tr } \rho = 1$ .

Show that the eigenvalues  $\lambda_i$  of  $\rho$  must satisfy  $0 \leq \lambda_i \leq 1$ .

By expressing the trace of  $\rho^2$  in terms of these eigenvalues, prove the inequality:

$$\text{Tr } \rho^2 \leq 1$$

Under what condition regarding the eigenvalues does  $\text{Tr } \rho^2 = 1$ ? (A state satisfying this condition is known as a *pure state*, whereas if  $\text{Tr } \rho^2 < 1$ , it is a *mixed state*. You'll see this topic in Part II Physics, if you do it.)

11 Let  $H$  be an  $n \times n$  Hermitian matrix representing the Hamiltonian (energy operator) of a quantum system. The time evolution of the state vector  $\boldsymbol{\psi}(t) = U(t)\boldsymbol{\psi}(0)$  is governed by the matrix exponential  $U(t) = e^{-iHt}$ .

Using the power series definition of the matrix exponential,

$$e^A = \sum_{n=0}^{\infty} \frac{A^n}{n!},$$

prove that  $U(t)$  is a unitary matrix.

If the initial state vector  $\boldsymbol{\psi}(0)$  is normalized such that  $|\boldsymbol{\psi}(0)|^2 = \boldsymbol{\psi}(0)^\dagger \boldsymbol{\psi}(0) = 1$ , show that the state vector remains normalized at all times  $t$ .