

Corrections to *ghm22@cam.ac.uk*. Star (★) indicates a harder question.

This course covers a bit of complex *analysis* as a precursor to complex *methods* (contour integrals) which come in Lent term.

Health warning: this is a difficult topic, and possibly whizzed through a bit in lectures. I've added a lot of additional questions here so you're comfortable with the full range of questions that might come up in the exam.

- 1 Define the radius of convergence of a power series of a complex variable  $z$ . Prove that the radius of convergence exists. Find the radii of convergence of the three series

$$\sum_{n=1}^{\infty} \frac{z^n}{n}, \quad \sum_{n=0}^{\infty} (\cosh n)z^n, \quad \text{and} \quad \sum_{n=0}^{\infty} [2^n + (-1)^n]z^n.$$

Show by example that a power series may or may not converge on its circle of convergence. Hence give an example of a series that is convergent but not absolutely convergent.

- 2 Calculate the Taylor series of the function  $f(z) = \log(1-z)$  about  $z=0$  and determine its radius of convergence. Now calculate the Taylor series for  $f(z)$  about  $z=i$  and determine the new radius of convergence. Comment.
- 3 Find the first three terms of the Taylor series expansion of the function:

$$f(z) = \frac{z}{e^z - 1}$$

about  $z=0$ . Show that the singularity at  $z=0$  is removable.

By identifying the positions of all other singularities of  $f(z)$  in the complex plane, determine the exact radius of convergence of this power series.

- 4 Where are the zeros and singularities of the following complex functions? Give the orders of the zeros, and classify the singularities.

$$\frac{(z-i)^2}{z+1}, \quad \frac{1}{1+z} - \frac{1}{1-z}, \quad \frac{1}{z^2+i}, \quad \sec^2 \pi z, \quad \sin z^{-2}, \quad \sinh \frac{z}{z^2-1}, \quad \frac{\tanh z}{z}.$$

- 5 To examine the behaviour of a complex function  $f(z)$  at infinity, we make the substitution  $z=1/\zeta$  and investigate the behaviour of the function  $g(\zeta) = f(1/\zeta)$  at  $\zeta=0$ .

Classify the behaviour at  $z=\infty$  (analytic, pole of a certain order, or essential singularity) for the following functions:

$$f(z) = z^4 - 3z^2 + 1, \quad f(z) = \frac{z^2 + 1}{2z^2 - z}, \quad f(z) = e^z, \quad f(z) = \sin\left(\frac{1}{z}\right).$$

- 6 What is meant by an *isolated singularity* of a complex function  $f(z)$ ?

Consider the function:

$$f(z) = \frac{1}{\sin(\pi/z)}.$$

Find all the singularities of  $f(z)$  in the finite complex plane. Show that the singularities at  $z_n = 1/n$  (where  $n \in \mathbb{Z}^+$ ) are isolated simple poles. Explain why the singularity at  $z=0$  is *not* an isolated singularity. Can  $f(z)$  be expanded in a Laurent series valid in a punctured disk  $0 < |z| < R$ ?

7 State the definition of complex differentiability for a function  $f(z)$  at a point  $z_0$ .

By considering paths parallel to the real and imaginary axes, derive the Cauchy–Riemann equations for  $f(z) = u(x, y) + iv(x, y)$ , where  $z = x + iy$ .

Show that the function  $f(z) = \bar{z}$  (the complex conjugate of  $z$ ) is nowhere complex differentiable. Show also that  $f(z) = |z|^2$  is complex differentiable at  $z = 0$ , but is not analytic there.

8 Show that if a function  $f(z) = u(x, y) + iv(x, y)$  is analytic in a domain  $\mathcal{D}$ , then both  $u(x, y)$  and  $v(x, y)$  satisfy Laplace’s equation:

$$\nabla^2 \phi = \frac{\partial^2 \phi}{\partial x^2} + \frac{\partial^2 \phi}{\partial y^2} = 0.$$

9 Show that the function on the plane defined by

$$\Phi(x, y) = \text{Im} \left\{ \frac{2}{\pi} \log \tanh(x + iy) \right\}$$

satisfies Laplace’s equation for  $x > 0$ . Show also that  $\Phi = 0$  on both  $y = 0$  and on  $y = \pi/2$  and that  $\Phi = 1$  on  $x = 0$  for  $0 < y < \pi/2$ . Deduce the steady-state temperature distribution in a semi-infinite two-dimensional bar of width  $L$ , with the (infinitely) long sides held at zero temperature and short side held at temperature  $T_0$ .

10 The real parts of three analytic functions  $f(x, y) = u(x, y) + iv(x, y)$  of  $z = x + iy$  are

$$\sin x \cosh y, \quad e^{y^2 - x^2} \cos 2xy, \quad \text{and} \quad \frac{x}{x^2 + y^2}$$

respectively. Use the Cauchy–Riemann equations to find their imaginary parts and hence deduce the forms of the complex functions.

Simplify your answers to become functions of  $z$  (rather than  $x$  and  $y$  separately). What do you notice about  $u(x, 0)$  in each case?

*Optional:* by considering  $df/dz = \partial f/\partial x$  and writing this in terms of  $u$  only, deduce a slightly simpler procedure to find the analytic function  $f$ .

11 Let  $f(z) = u(x, y) + iv(x, y)$  be an analytic function. Consider the geometric transformation from the  $xy$ -plane to the  $uv$ -plane defined by  $(x, y) \mapsto (u(x, y), v(x, y))$ . Find the Jacobian matrix  $J$  of this transformation. Use the Cauchy–Riemann equations to show that  $J$  can be expressed in the form

$$J = \begin{pmatrix} a & -b \\ b & a \end{pmatrix},$$

where  $a, b \in \mathbb{R}$ . Interpret what this matrix structure implies about the geometric interpretation of the local transformation.

Show that the families of level curves  $u(x, y) = a$  and  $v(x, y) = b$  (where  $a, b$  are constants) intersect orthogonally at all points where  $f'(z) \neq 0$ .

12 State the form the Laurent series for a function  $f$  with a pole of order  $n$  at the point  $z_0$ .

Consider the function

$$f(z) = \frac{1}{(z-1)(z-2)}.$$

Find the explicit Laurent series expansion of  $f(z)$  centred at  $z = 0$  valid for  $|z| < 1$ . Without calculating, how would you obtain the corresponding series valid for  $1 < |z| < 2$  and  $2 < |z|$ ?

- 13 Find the Laurent series expansion about  $z = 0$  for the function:

$$f(z) = z^3 \cos\left(\frac{1}{z}\right).$$

Determine the region of convergence for this series. Classify the singularity at  $z = 0$  and find the value of the coefficient of  $z^{-1}$ .

- 14★ The Casorati–Weierstrass theorem states that if  $z_0$  is an isolated essential singularity of  $f(z)$ , then in any neighbourhood of  $z_0$ , the function  $f(z)$  comes arbitrarily close to any complex number.

Illustrate this theorem using the function  $f(z) = e^{1/z}$  at  $z = 0$ . Given any non-zero complex number  $w = \rho e^{i\phi}$ , show that there is an infinite sequence of points  $z_n \rightarrow 0$  such that  $f(z_n) = w$ . What happens if  $w = 0$ ?